


Cell Reports

Cellular Mechanisms Underlying Behavioral State-Dependent Bidirectional Modulation of Motor Cortex Output

Graphical Abstract



Authors

Julia Schiemann, Paolo Puggioni, ..., Mark C.W. van Rossum, Ian Duguid

Correspondence

ian.duguid@ed.ac.uk

In Brief

Schiemann et al. show that, in mouse motor cortex, layer 5B pyramidal neuron firing rates are suppressed or enhanced during movement due to a global reduction in membrane potential variability and coincident noradrenaline-mediated depolarization in a subpopulation of neurons, respectively. Blocking noradrenergic input to M1 impairs motor coordination.

Highlights

- During movement, reduced V_m variance lowers spike probability in L5B output neurons
- Noradrenaline mediates a tonic depolarization in L5B_{enh} neurons during movement
- Noradrenaline selectively enhances signal-to-baseline ratio of L5B_{enh} neurons
- Blocking noradrenaline receptors in M1 reduces contralateral forepaw motor coordination



Cellular Mechanisms Underlying Behavioral State-Dependent Bidirectional Modulation of Motor Cortex Output

Julia Schiemann,^{1,4} Paolo Puggioni,^{1,2,3,4} Joshua Dacre,¹ Miha Pelko,^{2,3} Aleksander Domanski,¹ Mark C.W. van Rossum,² and Ian Duguid^{1,*}

¹Centre for Integrative Physiology and Patrick Wild Centre, University of Edinburgh, Hugh Robson Building, George Square, Edinburgh EH8 9XD, UK

²Institute for Adaptive and Neural Computation, School of Informatics, University of Edinburgh, Edinburgh EH8 9AB, UK

³Neuroinformatics Doctoral Training Centre, School of Informatics, University of Edinburgh, Edinburgh EH8 9AB, UK

⁴Co-first author

*Correspondence: ian.duguid@ed.ac.uk

<http://dx.doi.org/10.1016/j.celrep.2015.04.042>

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

SUMMARY


Neuronal activity in primary motor cortex (M1) correlates with behavioral state, but the cellular mechanisms underpinning behavioral state-dependent modulation of M1 output remain largely unresolved. Here, we performed *in vivo* patch-clamp recordings from layer 5B (L5B) pyramidal neurons in awake mice during quiet wakefulness and self-paced, voluntary movement. We show that L5B output neurons display bidirectional (i.e., enhanced or suppressed) firing rate changes during movement, mediated via two opposing subthreshold mechanisms: (1) a global decrease in membrane potential variability that reduced L5B firing rates (L5B_{suppressed} neurons), and (2) a coincident noradrenaline-mediated increase in excitatory drive to a subpopulation of L5B neurons (L5B_{enhanced} neurons) that elevated firing rates. Blocking noradrenergic receptors in forelimb M1 abolished the bidirectional modulation of M1 output during movement and selectively impaired contralateral forelimb motor coordination. Together, our results provide a mechanism for how noradrenergic neuromodulation and network-driven input changes bidirectionally modulate M1 output during motor behavior.

INTRODUCTION

Neuronal activity in layer 5 (L5) of primary motor cortex (M1) correlates with rhythmic voluntary movements (Armstrong and Drew, 1984a, 1984b). During walking or running, pyramidal neurons display changes in firing rate that reflect periods of coordinated muscle activity (Armstrong and Drew, 1984a; Beloozerova et al., 2003). Although spontaneous locomotor activity can be controlled by central pattern generators (CPGs) in the spinal

cord (Forssberg et al., 1980; Grillner, 1981; Grillner and Zangger, 1979), descending motor commands from M1 are integrated with ongoing rhythmic spinal cord signals and sensory input from the periphery to initiate, adjust, and maintain locomotor function (Armstrong and Drew, 1984a; Beloozerova et al., 2003; Orlovsky, 1972; Ueno and Yamashita, 2011). In lower mammals, such as cats, rabbits, and mice, discrete subpopulations of L5 output neurons display enhanced or suppressed (i.e., bidirectional) firing rate changes during locomotion (Armstrong and Drew, 1984a; Beloozerova et al., 2003; Costa et al., 2004). In rodents, these changes can be either abrupt, sustained changes—so-called on-off responses—or gradual frequency changes linked to the velocity of running (Costa et al., 2004). Although we are now beginning to understand how patterns of motor cortex activity relate to changes in behavioral state in rodents (i.e., quiet wakefulness to movement), the cellular mechanisms underpinning bidirectional modulation of M1 output during self-paced movement remain largely unresolved.

Several mechanisms could underlie the bidirectional modulation of M1 output, such as a change in cortical state-dependent network-driven input structure, intracortical or long-range excitatory input, and/or neuromodulation. During quiet wakefulness or slow-wave sleep, cortical networks remain in a synchronized state that consists of slow, large-amplitude oscillations in neuronal population activity (Cowan and Wilson, 1994; Steriade et al., 1993c). During active behavior, cortical networks enter an activated state characterized by a reduction in slow oscillations and, in some cases, an increase in higher frequency activity (Steriade et al., 1993b; Timofeev et al., 2001). This change profoundly alters the subthreshold V_m dynamics and spike output patterns of cortical pyramidal neurons (Castro-Alamancos, 2004; Castro-Alamancos and Oldford, 2002; Constantinople and Bruno, 2011; Crochet and Petersen, 2006). Thalamic activation promotes the cortical awake state and direct depolarization of superficial and deep-layer pyramidal neurons (Castro-Alamancos and Connors, 1996; Castro-Alamancos and Oldford, 2002; Constantinople and Bruno, 2013; Hirata and Castro-Alamancos, 2010; Poulet et al., 2012), suggesting input from the thalamus could contribute to changes in M1 output during motor



(legend on next page)

behavior. Anatomical and functional connectivity mapping have shown the presence of a strong top-down laminar organization of local excitatory microcircuits in M1, with feedforward projections from layer 2/3 (L2/3) targeting multiple classes of projection neurons in L5 (Kaneko et al., 1994; Weiler et al., 2008). Given that L2/3 pyramidal neurons can display dense clustered activity during head-restrained locomotion in mice (Dombeck et al., 2009), changes in descending excitation from upper-layer pyramidal neurons could be a mechanism for generating bidirectional modulation of M1 output. Alternatively, neuromodulators are important for cortical processing, with noradrenaline and acetylcholine release being associated with changes in arousal, vigilance, and behavioral state (Berridge and Waterhouse, 2003; Carter et al., 2010; Castro-Alamancos and Gulati, 2014; Constantinople and Bruno, 2011; Eggermann et al., 2014; Fu et al., 2014; Polack et al., 2013; Steriade et al., 1993a). Thus, how local, long-range, and neuromodulatory inputs regulate L5 pyramidal neuron V_m dynamics during changes in behavioral state remains to be fully established.

Here we combined in vivo patch-clamp recordings in awake mice with selective pharmacology to investigate the cellular mechanisms underpinning behavioral state-dependent modulation of motor cortex output. We found that changing behavioral state, from quiet wakefulness to movement, bidirectionally modulated (i.e., enhanced or suppressed) M1 output via two opposing subthreshold mechanisms: (1) a global decrease in network-driven, slow, large-amplitude V_m fluctuations, which reduced V_m variability, spike probability, and firing rates in L5B pyramidal neurons (L5B_{suppressed} neurons); and (2) a coincident increase in excitatory drive to a subpopulation of L5B neurons (L5B_{enhanced}), which depolarized mean V_m and enhanced firing rates. We found that the movement-related tonic depolarization in L5B_{enh} neurons was dependent on the interplay between ascending motor thalamic input, which maintained V_m near threshold, and noradrenergic input from the locus coeruleus (LC). The behavioral state-dependent release of noradrenaline increased the signal-to-baseline ratio (SBR) for movement-evoked responses in L5B_{enh} neurons. Selectively blocking noradrenergic input in the forelimb region of M1 significantly reduced motor coordination in the contralateral forelimb during motor behavior. Thus, our findings provide a mechanism for

how noradrenergic neuromodulation and network-driven input changes bidirectionally modulate M1 output during self-paced voluntary movement.

RESULTS

Membrane Potential Dynamics of L5B Pyramidal Neurons during Self-Paced, Voluntary Movement

To investigate the cellular mechanisms underpinning behavioral state-dependent modulation of M1 output, we obtained whole-cell patch-clamp recordings from L5B pyramidal neurons (forelimb motor cortex, 620–880 μm from the pial surface; see [Experimental Procedures](#); $n = 45$ neurons) during quiet wakefulness and self-paced, voluntary movements (i.e., walking, running, or grooming on a single-axis, cylindrical treadmill; [Figure 1A](#)). During periods of quiet wakefulness, all L5B pyramidal neurons displayed large-amplitude V_m fluctuations (V_m SD = 3.8 ± 0.2 mV) and a relatively depolarized average V_m ($V_m = -51.1 \pm 0.8$ mV). The interplay among mean V_m , distance from threshold, and V_m variability resulted in moderate basal firing rates (5.7 ± 0.6 Hz, range: 0.0–15.9 Hz; [Figures 1B–1K](#) and [S1](#)).

During switches in behavioral state (i.e., quiet wakefulness to movement), characterized by a low-amplitude, high-frequency local field potential signal in L5B ([Figure 1A](#)), the vast majority of L5B pyramidal neurons ($\sim 90\%$) displayed significant modulation of their basal firing rates. To functionally classify individual neurons, we compared the variability in quiet wakefulness firing rate with the average firing rate during self-paced movement (see [Experimental Procedures](#)). If the average movement-related firing rate was lower than the first percentile of the distribution of firing rate changes during quiet wakefulness, neurons were classified as suppressed (L5B_{supp}, $n = 17$; [Figures 1C](#) and [1F](#); [Table S1](#)), while neurons that displayed an average movement-related firing rate above the 99th percentile were classified as enhanced (L5B_{enh}, $n = 24$; [Figures 1E](#) and [1H](#); [Table S1](#)). A small proportion of L5B neurons ($n = 4/45$) did not significantly change their firing rates during movement and were classified as non-responding neurons (L5B_{n-r}; [Figures 1D](#) and [1G](#)). The proportion of L5 pyramidal neurons in which spike frequency decreased (38%), increased (53%), or did not change (9%) during movement was consistent with previous reports ([Beloozerova et al.](#),

Figure 1. Membrane Potential Dynamics of L5B Pyramidal Neurons in M1 during Self-Paced, Voluntary Movement

(A) Patch-clamp recording configuration in head-fixed mice mounted on a single-axis, cylindrical treadmill. Local field potential (LFP) recordings (black traces, $n = 2$ mice) of L5B activity and moderate speed (60 frames/s) digital imaging were used to confirm changes in behavioral state (quiet wakefulness to movement) and to calculate motion index (gray traces).
(B) Representative voltage traces from two L5B pyramidal neurons that displayed either a decrease (top) or increase (bottom) in firing rate during movement (light gray shading). The motion index (dark gray) defines the magnitude and duration of each forelimb movement. In this figure and all subsequent figures, action potentials have been truncated to highlight subthreshold V_m changes during movement.
(C–E) Representative change in firing rate probability distributions during quiet wakefulness (blue) and movement (gray) in L5B_{suppressed} (C), L5B_{non-responding} (D), and L5B_{enhanced} (E) neurons. Gray dotted lines represent the 1st (left) and 99th (right) percentiles. Solid colored lines represent the average firing rate change in L5B_{suppressed} (yellow), L5B_{non-responding} (black), and L5B_{enhanced} (purple) neurons during movement.
(F–H) Average firing rate during quiet wakefulness and movement in L5B_{suppressed} (F, $n = 17$), L5B_{non-responding} (G, $n = 4$), and L5B_{enhanced} (H, $n = 24$) neurons. Filled circles represent data from individual neurons. Insets depict the average firing rate \pm SEM during quiet wakefulness (Q) and movement (M). ** $p < 0.01$; ns, non-significant.
(I–K) Average V_m (left-hand plot) and V_m SD (right-hand plot) in L5B_{suppressed} (I, $n = 17$), L5B_{non-responding} (J, $n = 4$), and L5B_{enhanced} (K, $n = 24$) neurons during quiet wakefulness and movement. Solid gray lines represent data from individual neurons while open symbols represent mean \pm SEM. * $p < 0.05$, ** $p < 0.01$; ns, non-significant.

See also [Figures S1–S4](#) and [Tables S1](#) and [S2](#).




Figure 2. Input-Output Transformations in L5B_{supp} and L5B_{enh} Pyramidal Neurons during Movement

(A and B) Representative voltage traces (upper trace, black) during somatic EPSC-like current injections in vivo (lower trace, dark gray) in L5B_{supp} (A) and L5B_{enh} (B) pyramidal neurons during quiet wakefulness and movement (light gray shading).

(C and D) Input-output transformations in L5B_{supp} (C, n = 5) and L5B_{enh} (D, n = 6) neurons recorded in vivo during quiet wakefulness (blue) and movement (red). Symbols represent mean ± SD; solid lines are fits to a truncated error function.

(E and F) Mean change in spike probability for L5B_{supp} (E, n = 5) and L5B_{enh} (F, n = 6) neurons. Filled symbols represent data from individual neurons and black open squares represent mean ± SD.

See also Figure S2.

2003; Costa et al., 2004). Moreover, the functional classification of individual neurons remained consistent during repeated bouts of movement and was not dependent on the type of motor behavior being executed (Figure S1). To further demonstrate the coexistence of functionally distinct subpopulations of L5B pyramidal neurons in M1, we performed multiple recordings from the same mouse and identified L5B_{enh}, L5B_{supp}, and L5B_{n-r} pyramidal neurons during the execution of similar forelimb movements (i.e., repeated forepaw swing/stance cycles, n = 8 recordings from three mice; L5B_{supp}/L5B_{enh}/L5B_{n-r} ratio: 4.3:1, note similar ratio of functionally classified neurons when compared to the population data in Figure 1; Figure S1).

We next investigated the subthreshold mechanisms underpinning bidirectional modulation of M1 output. During movement, L5B_{supp} neurons displayed ~1 mV hyperpolarization in mean V_m ($p = 2 \times 10^{-2}$) and reduced V_m variability (V_m SD quiet = 3.5 ± 0.2 mV, V_m SD movement = 2.5 ± 0.1 mV, $p = 3 \times 10^{-4}$), which lowered the probability of reaching threshold and reduced overall firing rates (quiet 6.4 ± 1.0 Hz, movement 2.8 ± 0.6 Hz, $p = 3 \times 10^{-4}$; Figures 1F and 1I). In L5B_{enh} neurons, movement also reduced V_m variability (V_m SD quiet = 4.1 ± 0.3 mV, V_m SD

movement = 3.2 ± 0.2 mV, $p = 2 \times 10^{-3}$), but this was counteracted by a depolarization in average V_m (quiet -52.4 ± 1.1 mV, movement -47.9 ± 1.0 mV, $p = 2 \times 10^{-6}$), which significantly increased spike probability and firing rates (quiet 5.7 ± 0.8 Hz, movement 12.9 ± 1.5 Hz, $p = 2 \times 10^{-5}$; Figures 1H and 1K). Moreover, movement-related firing rate changes strongly correlated with the level of V_m depolarization in individual L5B_{enh} neurons (Figure S2). By contrast, V_m dynamics and firing rates of L5B_{n-r} neurons were not affected by the transition from quiet wakefulness to movement (Figures 1G and 1J). Interestingly, the functional classification of L5B pyramidal neurons (L5B_{supp} versus L5B_{enh}) was not dependent on their basic electrophysiological properties (Table S1) or the projection-class identity of individual neurons based on retrograde tracing and selective expression of the transcription factors CTIP2 (thick-tufted pyramidal tract [PT]-type neurons) and SATB2 (thin-tufted intratelencephalic [IT]-type neurons; Leone et al., 2008; Figures S3 and S4; Table S2). Together, our results suggest that movement-related modulation of L5B_{enh} firing rates is primarily mediated by a tonic depolarization in V_m , while reduced firing rates in L5B_{supp} neurons result from a moderate hyperpolarization and significant reduction in V_m variance.

L5B Input-Output Transformations during Voluntary Movement

Behavioral state-dependent changes in V_m dynamics can profoundly affect the integrative mode and output firing patterns of neocortical neurons. What effects do movement-related changes in V_m dynamics have on input-output transformations in M1 L5B pyramidal neurons? In principle, both changes in V_m SD and mean can influence the responsiveness and firing dynamics of a neuron (Chance et al., 2002; Hô and Destexhe, 2000). To test this, we performed current injection experiments (i.e., somatic injection of excitatory postsynaptic current [EPSC]-like waveforms) in a subset of L5B_{supp} and L5B_{enh} neurons in vivo (Figures 2A and 2B; Supplemental Experimental Procedures) and measured the spike probability during quiet wakefulness and voluntary movement. Although current injection at the soma disregards dendritic non-linearities, synaptic properties, and locations, it provides a robust measure to assess the relationship between synaptic conductances arriving at the soma and spike output probability during behavior. During movement L5B_{supp} neurons, which experience a decrease in V_m SD with relatively little change in mean V_m (Figure 1), displayed a 2-fold reduction in spike probability (Δ Spike probability = 0.6 ± 0.1 , n = 5; Figures 2C and 2E). By contrast, L5B_{enh} neurons, which experience a decrease in V_m SD and an increase in mean V_m (Figure 1), displayed a 2-fold increase in spike probability (Δ Spike probability = 1.7 ± 0.4 , n = 6; Figures 2D and 2F). Although both L5B subpopulations displayed moderate changes in input resistance during movement, they did not significantly differ from quiet wakefulness (n = 5 and 5, respectively, $p = 0.32$; Figure S2).

Changes in L5B Input Structure during Movement

To further investigate the mechanisms underpinning L5B_{supp} and L5B_{enh} neuron V_m dynamics, we explored changes in V_m spectral components before and after movement onset. During quiet wakefulness, we observed slow (1.5–4 Hz, δ frequency band), large-amplitude V_m fluctuations in all L5B pyramidal neurons




Figure 3. Movement Reduces Slow, Large-Amplitude V_m Fluctuations but Increases Excitatory Drive in $L5B_{enh}$ Neurons

(A and B) Representative high-time resolution voltage traces for $L5B_{supp}$ (A) and $L5B_{enh}$ (B) neurons during quiet wakefulness and movement (gray shading). (C and D) Low-time resolution wavelet spectrograms for $L5B_{supp}$ (C) and $L5B_{enh}$ (D) neurons during quiet wakefulness and movement. Representative examples correspond to neurons shown in (A and B). (E and F) Average V_m power density for $L5B_{supp}$ (E, $n = 17$) and $L5B_{enh}$ (F, $n = 24$) pyramidal neurons during quiet wakefulness (blue) and movement (red). Data represent mean \pm SD. Insets show average V_m power density between 5 and 40 Hz.

(Figures 3A–3D), which were suppressed during movement ($L5B_{supp}$ quiet $7.8 \pm 1.3 \text{ mV}^2$, movement $3.6 \pm 0.5 \text{ mV}^2$, $n = 17$, $p = 2 \times 10^{-3}$; $L5B_{enh}$ quiet $16.4 \pm 3.1 \text{ mV}^2$, movement $6.2 \pm 1.2 \text{ mV}^2$, $n = 24$, $p = 1 \times 10^{-4}$; Figures 3A–3H). The reduction in δ power led to reduced V_m SD, which together with a moderate hyperpolarization ($\sim 1 \text{ mV}$) could account for the reduction in spike probability observed in $L5B_{supp}$ pyramidal neurons during movement (Figures 1 and S2). In $L5B_{enh}$ neurons, the suppression of slow V_m fluctuations was counteracted by an increase in power (12–30 Hz) in the β frequency band (12–30 Hz; quiet $3.0 \pm 0.4 \text{ mV}^2$, movement $7.4 \pm 1.4 \text{ mV}^2$, $n = 24$, $p = 3 \times 10^{-5}$; Figures 3F and 3H). The magnitude of increased β power displayed a strong positive correlation with the magnitude of V_m depolarization in individual $L5B_{enh}$ neurons (Figure S2), suggesting this could be the source of the increased excitatory drive.

To examine this further, we developed an event detection algorithm to estimate the level of excitatory input during quiet wakefulness and movement. Due to the high frequency of afferent input (estimated range: 5–15 kHz, data not shown), we were unable to isolate single excitatory postsynaptic potentials (EPSPs). However, we could reliably detect compound synaptic inputs ($\geq 1 \text{ mV}$) occurring in a time window (5 ms) shorter than the average membrane time constant ($8.2 \pm 0.7 \text{ ms}$, $n = 10$; Figure S2). The detection threshold corresponded to twice the size of the average unitary synaptic response measured in L5 pyramidal neurons in vitro (Deuchars et al., 1994; Reyes and Sakmann, 1999). Events that occurred within $\pm 10 \text{ ms}$ of a spike were excluded from the analysis. During quiet wakefulness, we detected fast-rising compound EPSPs (range: 1–9.7 mV) with similar rates in both $L5B_{supp}$ and $L5B_{enh}$ pyramidal neurons (Figures 3I and 3J), indicating both subpopulations of neurons receive a comparable level of excitatory drive. During movement, the rate of compound events in $L5B_{supp}$ neurons was not affected ($n = 17$; Figure 3I), whereas $L5B_{enh}$ neurons displayed a significant increase in compound EPSP rate ($n = 24$; Figure 3J). Remarkably, we did not detect any compound events with amplitudes greater than 9.4 mV, even though neurons spent approximately 50% of the time $>10 \text{ mV}$ from threshold. Thus, $L5B_{enh}$ neurons appear to preferentially receive a net increase in excitatory drive during movement, which enhances the firing rate by depolarizing mean V_m and increasing spike probability.

Effects of Local and Long-Range Input to $L5B$ Pyramidal Neurons during Self-Paced Movement

To investigate the possible source(s) of the increased excitatory drive to $L5B_{enh}$ neurons, we examined the activity of local and long-range inputs from L2/3 and motor thalamus, respectively.

(G and H) Average V_m power in δ (1.5–4 Hz) and β (12–30 Hz) frequency bands in $L5B_{supp}$ (G, $n = 17$) and $L5B_{enh}$ (H, $n = 24$) pyramidal neurons during quiet wakefulness and movement. Gray lines represent data from individual neurons and black symbols represent mean \pm SEM. $**p < 0.01$.

(I and J) Average rate density of compound synaptic events in $L5B_{supp}$ (I, $n = 17$) and $L5B_{enh}$ (J, $n = 24$) pyramidal neurons during quiet wakefulness (blue) and movement (red). Data represent mean \pm SD. $**p < 0.01$; ns, non-significant.

See also Figure S2.




Figure 4. Effect of Descending L2/3 and Ascending Motor Thalamic Input on L5B Pyramidal Neuron V_m Dynamics during Quiet Wakefulness and Movement

(A) Representative voltage trace shows an L2/3 pyramidal neuron during quiet wakefulness and movement (gray shading). (B–D) Average firing rate (B), mean V_m (C), and V_m SD (D) in L2/3 pyramidal neurons (gray symbols, $n = 8$) before and after movement. Filled circles represent data from individual neurons while square symbols represent mean \pm SEM. Inset in (B) depicts average L2/3 pyramidal neuron firing rate during quiet wakefulness (Q) and movement (M). ns, non-significant.

Previous studies have shown that M1 L2/3 neurons can be highly active during head-restrained locomotion on a spherical treadmill (Dombeck et al., 2009) and this descending excitation could potentially influence the activity of output neurons in L5 (Weiler et al., 2008). To test this possibility, we obtained whole-cell patch-clamp recordings from L2/3 pyramidal neurons (180–420 μ m from the pial surface; Figure 4A). During quiet wakefulness, L2/3 neurons displayed relatively low firing rates, which were unaffected by the onset of movement (quiet 0.6 ± 0.3 Hz, movement 0.6 ± 0.4 Hz, $n = 8$; Figure 4B). Although the average V_m of L2/3 neurons depolarized by ~ 4 mV (Figure 4C), this was counteracted by a significant reduction in V_m SD (Figure 4D), which maintained baseline spike probability and firing rates. Given that our sample of L2/3 neurons displayed low spike rates during both quiet wakefulness and movement, this suggests that descending input from L2/3 is unlikely to be the primary source of the increased excitatory drive to L5B_{enh} neurons in our experimental paradigm (see also Dombeck et al., 2009 and Discussion).

Given that thalamocortical neuron firing rates vary markedly depending on behavioral state and ventroanterior/ventrolateral (VA/VL) thalamic input to M1 displays bidirectional modulation during simple locomotion (Marlinski et al., 2012), we investigated the role of motor thalamus in regulating M1 output during movement. Blocking thalamic input by local infusion of the GABA_A receptor agonist muscimol into the VA/VL complex (Experimental Procedures; Figure 4E) enhanced the amplitude of slow, large-amplitude V_m fluctuations (control V_m SD = 3.8 ± 0.2 mV versus thalamic inactivation SD = 5.1 ± 0.6 mV; $n = 45$ and $n = 6$, respectively; $p = 3.4 \times 10^{-2}$, Mann-Whitney U test) and produced a hyperpolarizing shift in average V_m (control $V_m = -51.1 \pm 0.8$ mV versus thalamic inactivation $V_m = -62.5 \pm 3.6$ mV; $n = 45$ and $n = 6$, respectively; $p = 8.0 \times 10^{-4}$), which significantly reduced the basal firing rate of L5B neurons compared to control conditions (control 5.7 ± 0.6 Hz versus thalamic inactivation 1.1 ± 0.5 Hz; $n = 45$ and $n = 6$, respectively; $p = 1.1 \times 10^{-3}$; Figures 4G and 4H; see also Figure 1). The hyperpolarization associated with thalamic inactivation increased the distance to threshold

(E) Schematic representation shows an L5B pyramidal neuron recording after inactivation of ipsilateral motor thalamus (MTh) by local perfusion of muscimol. (F) Representative voltage trace showing an L5B pyramidal neuron after ipsilateral inactivation of motor thalamus.

(G and H) Average firing rate (G) and mean V_m (H) in L5B pyramidal neurons after motor thalamic inactivation ($n = 6$). Filled circles represent data from individual neurons while the square symbol in (H) represents the mean \pm SEM. Inset in (G) depicts the average firing rate of L5B neurons during quiet wakefulness (Q) and movement (M). ns, non-significant.

(I) Change in average V_m (ΔV_m) during movement in the presence (Ctrl, open symbols, $n = 41$) and absence of motor thalamic input (gray symbols, $n = 6$). Control data (Ctrl) were taken from Figure 1 for comparison. Mann-Whitney U test; ns, non-significant.

(J) Probability density distributions of ΔV_m variability across the L5B pyramidal neuron population (Ctrl and MTh inact.), measured as the SD of the ΔV_m distributions shown in (I) (Population ΔV_m SD) using bootstrap analysis (10,000 bootstrap replicates). Black dashed line represents population ΔV_m variability distribution in control (Ctrl), and gray shading represents population ΔV_m variability distribution following motor thalamic inactivation. Control data (Ctrl) were taken from Figure 1 for comparison. F test; ns, non-significant. See also Figure S5.

(data not shown) such that movement-related firing rate changes were abolished (Figure 4G), precluding the functional classification of L5B_{supp} and L5B_{enh} neurons. However, during movement 50% of L5B neurons ($n = 3/6$) still experienced a 5–10 mV depolarization in mean V_m (Figure 4I) and increased rate of compound EPSPs (Figure S5), similar to that observed in L5B_{enh} neurons under control conditions (Figures 1 and 3). We analyzed this further by plotting the ΔV_m variability across the L5B pyramidal neuron population, measured as the SD of the ΔV_m distributions shown in Figure 4I (population ΔV_m SD), using bootstrap analysis (10,000 bootstrap replicates; Figure 4J). We found that motor thalamic inactivation did not affect the population ΔV_m variability in L5B pyramidal neurons compared to control (Figures 4I and 4J), suggesting input from the motor thalamus—either direct or indirect—is essential for maintaining L5B pyramidal neuron V_m near threshold, but is unlikely to be the main source of the increased excitatory drive.

Noradrenergic Neuromodulation Selectively Enhances Excitatory Drive and Signal-to-Baseline Ratio in L5B_{enh} Neurons

Given that the movement-related increase in excitatory drive and tonic depolarization in L5B_{enh} neurons could not be directly explained by increased excitation from L2/3 or motor thalamus, we next explored the role of noradrenergic neuromodulation, which has been shown to be important during changes in arousal, attention, and behavioral state (Berridge and Waterhouse, 2003; Carter et al., 2010; Castro-Alamancos and Gulati, 2014; Constantinople and Bruno, 2011; Polack et al., 2013). Selective immunohistochemical staining for the noradrenaline transporter (NAT), expressed exclusively in noradrenergic axons (Lorang et al., 1994), revealed dense axonal innervation of all layers in forelimb M1 (Figure 5A). To test the importance of noradrenergic input in regulating L5B pyramidal neuron V_m dynamics during movement, we topically applied α_1 , α_2 , and β noradrenergic receptor antagonists (1 mM prazosin, 1 mM yohimbine, and 1 mM propranolol, respectively) to the forelimb region of M1 (Figures 5B and 5C). The local infusion of noradrenergic receptor antagonists via the craniotomy selectively disrupted noradrenergic signaling in forelimb M1 (Figure S6), whereas direct manipulation of LC activity, via electrical stimulation or optogenetics, would have widespread effects across many brain areas and spinal cord circuits. Moreover, topical application was preferred due to the technical limitations of simultaneously pressure ejecting drugs at multiple sites along the entire somatodendritic length of L5B pyramidal neurons during intracellular recordings.

Blocking noradrenergic receptors reduced the mean V_m (control -51.1 ± 0.8 mV versus noradrenergic receptor blockade -56.6 ± 1.6 mV; $p = 4.0 \times 10^{-3}$) and quiet wakefulness firing rate of L5B pyramidal neurons (control 5.7 ± 0.6 Hz versus noradrenergic receptor blockade 1.9 ± 0.3 Hz; $n = 45$ and $n = 16$, respectively; $p < 1 \times 10^{-4}$; Figures 5D and 5E), and significantly reduced the proportion of L5B neurons that displayed enhanced firing rates during movement (control L5B_{enh} 24/45 neurons [53.3%] versus noradrenergic receptor blockade L5B_{enh} 2/16 neurons [12.5%]; $p < 1 \times 10^{-2}$; Figures 5F and 5G). The change in relative distribution of L5B_{supp}/L5B_{enh} neurons could be ex-

plained in part by the moderate hyperpolarization in V_m and increased distance to threshold during movement (Figure S6). Although noradrenergic receptor blockade did not affect the mean population ΔV_m compared to control conditions, due to both distributions being centered around 0 mV (Figure 5H), we did observe a significant decrease in ΔV_m variability across the L5B pyramidal neuron population, measured as the SD of the ΔV_m distributions shown in Figure 5H (population ΔV_m SD) using bootstrap analysis (10,000 bootstrap replicates; Figure 5I). Consistent with the idea that noradrenergic signaling underpins a large proportion of the increased excitatory drive to L5B_{enh} neurons during movement, blocking noradrenergic receptors also abolished the increase in V_m β -band power (Figure 5J) and rate of compound synaptic events associated with movement (Figures 5K and S5).

Given that pre-application of noradrenergic receptor antagonists precludes the prior identification of L5B_{enh} neurons prior to receptor blockade, we also performed long-term (40- to 80-min) recordings from identified L5B_{enh} neurons before (Figure 6A) and after (Figure 6B) receptor block. If noradrenergic neuromodulation underpins the V_m depolarization in L5B_{enh} neurons during movement, then blocking noradrenergic receptors should have a disproportionately larger effect on movement-related firing rates compared to quiet firing rates. Accordingly, we found that receptor blockade resulted in a modest, time-dependent reduction in L5B_{enh} basal firing rates and a strong suppression of movement-related firing (Figure 6C). The drug diffusion and time dependency of the antagonist effects in L5B were consistent with our dye diffusion mapping results (Figure S6). To assess the extent to which noradrenaline facilitates L5B_{enh} output during movement, we examined the Signal-to-Baseline Ratio (SBR), defined as the ratio of the movement-related spike rate to the spike rate during quiet wakefulness. Blocking noradrenergic neurotransmission significantly reduced the SBR compared to control conditions (sham control SBR: 1.1 ± 0.1 , noradrenergic receptor antagonist SBR: 0.3 ± 0.1 ; $p = 6 \times 10^{-3}$; $n = 3$ and 3 , respectively; Figure 6D).

Since descending M1 output is essential for maintaining normal locomotor function (Armstrong and Drew, 1984a; Beloozerova et al., 2003; Orlovsky, 1972; Ueno and Yamashita, 2011), we investigated whether there was a behavioral correlate of reduced M1 output during noradrenergic receptor blockade by conducting a series of behavioral experiments using head-restrained mice habituated to walk/run on a cylindrical runged treadmill (Figure 6E). This experimental paradigm facilitates the analysis of precise forepaw placements during locomotion, which was not possible on the conventional single-axis cylindrical treadmill shown in Figure 1A. Although classified as complex locomotion, this paradigm generates only subtle differences in forelimb muscle activity/wrist movements and comparable changes in M1 activity when compared to simple locomotion on a linear treadmill (Beloozerova et al., 2010; Marlinski et al., 2012). Selectively blocking noradrenergic receptors in forelimb M1 significantly decreased the number of precise contralateral forepaw placements compared to sham controls (precise forepaw placements 60 min after: sham saline $86.8\% \pm 0.7\%$, noradrenergic receptor antagonists $70.5\% \pm 1.7\%$; $n = 3$ and 5 , respectively; $p < 4.0 \times 10^{-4}$; Figure 6F) or ipsilateral forepaw placements




Figure 5. Blocking Input from the LC Reduces Behavioral State-Dependent Increase in Excitatory Drive in L5B_{enh} Neurons

(A) Noradrenergic axons in the forelimb region of M1 were labeled using an anti-noradrenaline transporter antibody and secondary antibody conjugated to AlexaFluor 488.

(B) Schematic representation of an L5B pyramidal neuron recording after blocking noradrenergic input from the LC.

(C) Representative voltage trace from a L5B pyramidal neuron in the absence of noradrenergic input.

(D and E) Average V_m (D) and firing rate (E) of L5B pyramidal neurons during quiet wakefulness in the absence (open symbols, n = 45) and presence (green symbols, n = 16) of noradrenergic receptor (NA-R) antagonists. Filled circles represent data from individual neurons, black bars represent mean \pm SEM. Control data (Ctrl) were taken from the dataset presented in Figure 1 for comparison. Mann-Whitney U test, *p < 0.017, **p < 0.003.

(F) Average firing rate of L5B pyramidal neurons in the presence of noradrenergic receptor antagonists (n = 16) during quiet wakefulness and movement. Filled circles represent data from individual neurons.

(G) Relative distributions of L5B_{supp}, L5B_{enh}, and L5B_{n-r} neurons in the absence (Ctrl) and presence (NA-R antagonists, n = 16) of noradrenergic receptor antagonists. Control data (Ctrl) were taken from the dataset presented in Figure 1 for comparison. Chi-square test, **p < 0.01.

(H) Change in average V_m (ΔV_m) during movement in the absence (Ctrl, n = 41) and presence of noradrenergic receptor antagonists (green symbols, n = 16). Control data (Ctrl) were taken from Figure 1 for comparison. Mann-Whitney U test; ns, non-significant.

(I) Probability density distributions of ΔV_m variability across the L5B pyramidal neuron population (Ctrl and NA-R antagonists), measured as the SD of the ΔV_m distributions shown in (H) (Population ΔV_m SD) using bootstrap analysis (10,000 bootstrap replicates). Black dashed line represents population ΔV_m variability distribution in control (Ctrl), and green shading represents population ΔV_m variability distribution following noradrenergic receptor blockade. Control data (Ctrl) were taken from Figure 1 for comparison. F test, *p < 0.025.

(J) Average L5B pyramidal neuron V_m power in the β frequency band (12–30 Hz) during quiet wakefulness and movement in the presence (Ctrl: L5B_{supp}, n = 17 and L5B_{enh}, n = 24) and absence of noradrenergic input (NA-R antagonists, n = 16). Solid lines represent data from individual neurons, symbols represent mean \pm SEM. Control data (Ctrl) were taken from Figure 3 for comparison. *p < 0.05.

(K) Average rate density of compound synaptic events in L5B pyramidal neurons during quiet wakefulness (blue) and movement (red) in the absence of noradrenergic input (n = 16). Compare with Figures 3I and 3J.

See also Figures S5 and S6.

(data not shown). Together, our results demonstrate that noradrenergic input from the LC is necessary for controlling M1 output and motor coordination during self-paced voluntary movement.

DISCUSSION

In this paper we present three main findings. First, we show that behavioral state-dependent bidirectional modulation of M1 output is governed by two opposing subthreshold mechanisms (1) a global decrease in network-driven, slow, large-amplitude V_m fluctuations, which reduced V_m variability, spike probability, and firing rates in L5B_{supp} neurons; and (2) a coincident increase in excitatory drive in a subpopulation of L5B neurons (L5B_{enh}), which increased spike probability and firing rates. Second, we demonstrate that the movement-related tonic depolarization in L5B_{enh} neurons requires the interplay between ascending input from the motor thalamus, which maintained V_m near threshold, and noradrenergic input from the LC, which enhanced the SBR for movement-evoked responses. Finally, we show that selective




Figure 6. Blocking Noradrenergic Input Reduces the SBR in L5B_{enh} Neurons and Impairs Contralateral Forepaw Motor Coordination

(A and B) Schematic representation of the experimental design and representative voltage traces from an L5B_{enh} pyramidal neuron prior to (A) and after (B, >30 min) topical application of noradrenergic receptor antagonists. Gray shading depicts movement.

(C) Time course shows quiet (blue) and movement-related (red) firing rates in an L5B_{enh} pyramidal neuron before and after noradrenergic receptor blockade (green bar).

(D) Movement-induced SBR in L5B_{enh} pyramidal neurons before and >30 min after application of noradrenergic receptor antagonists (green circles, n = 3) or saline (gray circles, n = 3). Square symbols represent mean ± SEM. Unpaired t test, **p < 0.01.

(E) Behavioral assessment of forepaw placement precision in head-fixed mice mounted on a single-axis, cylindrical runged treadmill. Video sequences were used to score contralateral and ipsilateral forepaw placements.

(F) Percentage of precise contralateral forepaw placements before and 60 min after application of noradrenergic receptor antagonists (green circles, n = 5) or saline (gray circles, n = 3). Square symbols represent mean ± SEM. Unpaired t test, **p < 0.01.

See also Figure S6.

blockade of noradrenaline signaling in forelimb M1 reduces motor coordination in the contralateral forelimb, resulting in a significant decrease in precise forepaw placements. Together, our findings reveal the subthreshold and circuit mechanisms that regulate behavioral state-dependent bidirectional modulation of M1 output during self-paced, voluntary movement.

Behavioral State-Dependent Modulation of Input-Output Transformations in L5

Physiologically relevant changes in V_m variance or mean have been shown to profoundly influence neuronal input-output transformations (Chance et al., 2002; Hó and Destexhe, 2000). But, this has never been explored in L5 pyramidal neurons in the awake cortex. Our current injection experiments in vivo demonstrate that changes in V_m SD (L5B_{supp}) or V_m SD and mean (L5B_{enh}) have quantitatively similar—but functionally opposing—effects on spike probability when examined over a behaviorally relevant input amplitude range (1–10 mV). This similarity arises due to the non-linear relationship between V_m and firing probability, such that moderate depolarization can produce a non-linear additive increase in the sensitivity of a neuron to small-amplitude inputs, while decreased V_m SD produces a divisive reduction in input sensitivity (Brozović et al., 2008; Murphy and Miller, 2003). The behavioral state-dependent bidirectional modulation of neuronal responsiveness in L5B pyramidal neurons (i.e., increased or decreased spike probability) could facilitate the routing of sensorimotor information through specific M1 neuronal assemblies during movement.

Local and Long-Range Inputs to M1 during Self-Paced Voluntary Movement

M1 receives input from a variety of brain areas (e.g., ipsilateral primary and secondary somatosensory cortices, secondary motor cortex, and orbitofrontal cortex), with ascending input from motor thalamus and descending input from L2/3 providing strong feedforward excitation directly to L5B neurons (Castro-Alamancos and Connors, 1996; Hooks et al., 2013; Weiler et al., 2008). We found that movement did not affect firing rates in our sample of L2/3 pyramidal neurons, suggesting that descending excitation from L2/3 may not be the primary source of the tonic depolarization in L5B_{enh} neurons during simple locomotion on a cylindrical treadmill. These findings are in direct contrast to a previous study by Dombeck and colleagues, where locomotion on a spherical treadmill resulted in large-scale, clustered activity of L2/3 neurons in mouse forelimb M1 (Dombeck et al., 2009). The reason for this discrepancy is unclear. One possibility is that our recordings undersampled L2/3 population activity; however, if locomotion induced dense activity similar to that observed in Dombeck et al., (2009), we would have expected to observe movement-related firing rate changes in a significant proportion of our intracellular recordings. Moreover, Dombeck and colleagues did not identify individual neuronal subtypes, so the large-scale activity could be due, in part, to elevated L2/3 interneuron activity. Alternatively, dense L2/3 activity could result from mice having to balance and oppose the inertial forces of a rotating air-cushioned ball when changing direction. In principle, this could generate a sensorimotor

mismatch between the rotational direction of the ball and the intended movement trajectory of the mouse, leading to continuous sensory feedback to M1. Thus, it will be important for future studies to investigate the extent to which descending L2/3 input contributes to L5B frequency modulation during simple versus complex motor behaviors.

Direct thalamic input to cortical pyramidal neurons can drive output by reducing slow V_m fluctuations, depolarizing mean V_m , and reducing the distance to threshold (Castro-Alamancos and Connors, 1996; Constantinople and Bruno, 2013; Hirata and Castro-Alamancos, 2010; Poulet et al., 2012). Consistent with previous findings in sensory cortex, we found that inactivation of thalamus (VA/VL region) increased slow, large-amplitude V_m fluctuations, but did not abolish the activated state during behavior (Hirata and Castro-Alamancos, 2010; Poulet et al., 2012), suggesting thalamic input to M1 is sufficient but not necessary for generating the activated cortical state. However, ascending motor thalamic input—direct or indirect—appears to be necessary for maintaining the average V_m relatively close to threshold, providing a mechanism whereby subtle changes in input structure can generate positive or negative changes in M1 output during movement.

Noradrenergic Neuromodulation

We have shown that noradrenaline release during different behavioral states (i.e., quiet wakefulness versus movement) has profound effects on M1 cortical dynamics. Similar to thalamic inactivation, blocking noradrenergic input from the LC reduced basal firing rates by hyperpolarizing mean V_m and increasing distance to threshold, suggesting tonic input from both the LC and motor thalamus are necessary to generate moderate firing rates in L5B pyramidal neurons during quiet wakefulness. Our finding that noradrenaline generated a tonic depolarization in a selected subpopulation of L5B pyramidal neurons differs from results obtained in superficial layers of sensory cortex (Polack et al., 2013), highlighting the importance of understanding the sublayer-specific effects of noradrenaline in the awake cortex. In primary visual cortex (V1), locomotion-dependent noradrenaline release generates a global depolarization of L2/3 pyramidal neurons, which may enhance visual attention by increasing the gain and signal-to-noise ratio of visually evoked responses (Bennett et al., 2013; Polack et al., 2013).

The fact that we also observed a movement-related tonic depolarization in the majority of M1 L2/3 pyramidal neurons, which was abolished by noradrenergic receptor blockade (Figure S6), suggests that noradrenaline may differentially affect cortical processing in superficial versus deep-layer pyramidal neurons during active behavior. Topical application of high concentrations of noradrenergic receptor antagonists could potentially produce off-target effects. However, given that low doses of antagonists affect L2/3 V_m dynamics in the same way as high concentrations, albeit smaller in magnitude, suggests relatively selective antagonist effects (Polack et al., 2013). Although noradrenaline appears to underpin the majority of the locomotion-dependent V_m depolarization in V1, cholinergic disinhibition of somatostatin-containing interneurons is likely to further enhance behavioral state-dependent gain modulation (Fu et al., 2014). We did not directly test the role of acetylcholine in our study, but given

its importance in regulating V_m dynamics in other cortical areas (Constantinople and Bruno, 2011; Eggermann et al., 2014; Favero et al., 2012; Fu et al., 2014; Polack et al., 2013), it will be important for future studies to investigate its role in M1 during motor behavior.

How does noradrenaline generate the tonic depolarization in L5B_{enh} neurons during movement? Previous studies have shown that noradrenaline modulates voltage-dependent and voltage-independent potassium conductances and hyperpolarization-activated cyclic nucleotide-gated (HCN) channels, thus generating a tonic depolarization by reducing the spike after-hyperpolarization and prolonging the depolarizing effect of excitatory synaptic inputs (Favero et al., 2012; Sheets et al., 2011; Wang et al., 2007; Wang and McCormick, 1993). This combined with modulation of basal firing rates is thought to alter the signal-to-noise ratio of neuronal responses to synaptic input (Berridge and Waterhouse, 2003). Alternatively, we cannot rule out the possibility that noradrenaline selectively reduces the activity of local GABAergic interneurons, thus releasing L5B_{enh} neurons from inhibition and generating a depolarization in V_m . Therefore, identifying the specific expression patterns and subcellular localization of α and β adrenergic receptors in excitatory and inhibitory neurons in M1 will be an important next step in understanding how noradrenaline exerts its sublayer- and cell-type specific effects.

Functional Implications

What function does behavioral state-dependent bidirectional modulation of L5 output serve? The flexible modulation of L5B output channels (PT type and IT type) provides an important control mechanism to modulate and update activity patterns in downstream cortical and subcortical areas during changes in behavioral state. PT output provides online information about the state of cortical activation to downstream areas involved in motor control. This continuously updating flow of information generates a basic pattern of input to brainstem and spinal cord circuits in order to generate appropriate behavioral responses in accordance with changes in behavioral state. We demonstrate that blocking noradrenergic receptors in forelimb M1 selectively disrupts motor coordination in the contralateral forepaw, thus confirming the importance of noradrenergic neuromodulation and descending M1 output for motor control. Given that output from sensory and non-sensory cortices have overlapping downstream targets (Hattox and Nelson, 2007; Kita and Kita, 2012), we speculate that our findings might generalize to other cortical output layers and that noradrenergic neuromodulation and network-driven input changes are common mechanisms to bidirectionally modulate cortical output during active behavior.

EXPERIMENTAL PROCEDURES

Animals and Surgery

All experiments and procedures involving animals were approved by the University of Edinburgh local ethical review committee and performed under license from the UK Home Office in accordance with the Animal (Scientific Procedures) Act 1986. Male C57BL/6 mice (5–12 weeks old, 20–25 g, two to six animals per cage, maintained on a reversed 12:12-hr light:dark cycle with ad libitum access to food and water) were implanted with a small lightweight headplate (0.75 g) using cyano-acrylate glue and dental acrylic. All surgical

procedures were performed under 1.5% isoflurane anesthesia. After 24- to 48-hr recovery, a craniotomy (300 × 300 μ m) was performed and the dura removed above the right forelimb region of M1. Using intracortical microstimulation (see [Supplemental Experimental Procedures](#)), the center of M1_{FL} was located 0.7 mm rostral and 1.5 mm lateral to bregma. The craniotomy was sealed with (1.5%) agar and Kwik-Cast sealant (WPI) and mice recovered for 2 hr before recording commenced.

Motion Index and Motor Pattern Discrimination

An optical encoder was used to capture movement of the treadmill and locomotion was defined as periods with speed > 0.01 m/s for more than 2 s. Changes in behavioral state (quiet wakefulness to movement [grooming or locomotion]) were captured using an elevated, front-mounted, moderate-speed (60 frames/s) digital video camera synchronized with each electrophysiological recording.

In Vivo Electrophysiology and Pharmacology

Mice were habituated to the head restraint and experimental setup for 45–60 min before each recording session. Whole-cell patch-clamp recordings were obtained from awake head-restrained mice at a depth of 180–420 μ m (layer 2/3) or 620–880 μ m (layer 5B) from the pial surface, using a Multiclamp 700B amplifier (Molecular Devices). The signal was filtered at 10 kHz and acquired at 20 kHz using PClamp 10 software in conjunction with a DigiData 1440 DAC interface (Molecular Devices). No bias current was injected during recordings and the membrane potential was not corrected for junction potential. Resting membrane potentials were recorded immediately after attaining the whole-cell configuration (break-in). Series resistances (R_s) ranged from 15 to 40 M Ω and experiments were terminated if R_s exceeded 60 M Ω . Current injection was performed only if R_s < 35 M Ω . Patch pipettes (5–7 M Ω) were filled with internal solution (285–295 mOsm) containing: 135 mM K-gluconate, 4 mM KCl, 10 mM HEPES, 10 mM sodium phosphocreatine, 2 mM MgATP, 2 mM Na₂ATP, 0.5 mM Na₂GTP, and 2 mg/ml biocytin (pH adjusted to 7.2 with KOH). External solution contained: 150 mM NaCl, 2.5 mM KCl, 10 mM HEPES, 1 mM CaCl₂, and 1 mM MgCl₂ (adjusted to pH 7.3 with NaOH).

For inactivation of the motor thalamus, the GABA_A receptor agonist muscimol (1 mM muscimol hydrobromide, Sigma-Aldrich) was dissolved in external solution, and 100 nl was stereotactically injected into the right VA/VL complex (−1 mm caudal, 1.1 mm lateral to bregma, and 3.2 mm below the pial surface). Whole-cell patch-clamp recordings of L5B pyramidal neurons were carried out approximately 2 hr after muscimol injection.

To block noradrenergic receptors, a mixture of α 1, α 2, and β noradrenergic receptor antagonists (1 mM prazosin, yohimbine, and propranolol; Sigma-Aldrich) in external solution (adjusted to pH 7.3) was applied topically to the craniotomy and recordings were performed >40 min after antagonist application.

Functional Classification of Recorded Neurons

For each L5B cell, we (1) divided quiet periods into 1-s epochs; (2) randomly assigned each epoch into two groups, quiet 1 (q1) and quiet 2 (q2); and (3) calculated the firing rate difference between q1 and q2. We repeated steps (1) to (3) 10,000 times for each cell to obtain the distribution probability of the difference of firing rate in q1 and q2 (see [Figures 1C–1E](#)). If during movement the firing rate change was higher than the 99th percentile or lower than the 1st percentile, we classified the neuron as enhanced or suppressed, respectively. If the firing rate change fell within the first and 99th percentiles, the cell was classified as non-responding.

Statistical Analyses

Summary data are expressed as mean \pm SEM unless otherwise stated. Statistical significance was determined using Wilcoxon signed-rank tests (paired data) and rank-sum tests (unpaired data) unless otherwise stated. Wilcoxon signed-rank tests on the areas underlying the rate-density curves were used in [Figures 3I](#) and [3J](#). The relative distribution of functional phenotypes (L5B_{supp}, L5B_{enh}, and L5B_{nr}) was analyzed using Pearson chi-square test statistics (based on 10⁵ permutations). Statistical significance in population ΔV_m variability ([Figures 4J](#) and [5I](#)) was determined using two-sample F tests. To de-

pict the variance of the underlying populations, 10,000 bootstrap samples (random samples with replacement) of each population were taken, and a probability density function of the variances of the bootstrap samples was plotted. For statistical tests, $p < 0.05$ was considered significant (* $p < 0.05$ and ** $p < 0.01$). For repeated statistical comparisons with the control dataset, resulting p values were compared to Bonferroni-corrected alpha levels and stated accordingly.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, six figures, and two tables and can be found with this article online at <http://dx.doi.org/10.1016/j.celrep.2015.04.042>.

AUTHOR CONTRIBUTIONS

J.S., P.P., M.C.W.v.R., and I.D. designed the experiments. All experiments were carried out by J.S., P.P., and J.D. with help from I.D. Analysis was performed by P.P., J.S., M.P., and J.D. A.D. wrote MATLAB scripts to extract forepaw motion statistics. Neuronal reconstructions were carried out by J.S., P.P., J.D., and I.D. I.D., P.P., and J.S. wrote the manuscript, and all authors contributed to discussion and interpretation of the results.

ACKNOWLEDGMENTS

We are grateful to M. Nolan, M. London, P. Magill, K. Harris, T. Branco, L. Acerbi, M. Ludwig, N. Rochefort, and members of the I.D. lab for helpful discussions and for comments on the manuscript; M. Cowper-Coles for neuronal reconstructions; and T. Theil for advice on CTIP2/SATB2 immunohistochemistry. Confocal microscopy was performed in the IMPACT Imaging Facility at the University of Edinburgh. This work was supported by grants from the Engineering and Physical Sciences Research Council, Eurospin Erasmus Mundus Program, Leopoldina fellowship program (LPDS 2012-11 to J.S.), the Shirley Foundation, Medical Research Council (MR/K014137/1, awarded to P. Kind), and a Wellcome Trust Career Development fellowship (WT086602MF) to I.D.

Received: June 11, 2014

Revised: March 16, 2015

Accepted: April 20, 2015

Published: May 14, 2015

REFERENCES

- Armstrong, D.M., and Drew, T. (1984a). Discharges of pyramidal tract and other motor cortical neurones during locomotion in the cat. *J. Physiol.* 346, 471–495.
- Armstrong, D.M., and Drew, T. (1984b). Locomotor-related neuronal discharges in cat motor cortex compared with peripheral receptive fields and evoked movements. *J. Physiol.* 346, 497–517.
- Beloozerova, I.N., Sirota, M.G., and Swadlow, H.A. (2003). Activity of different classes of neurons of the motor cortex during locomotion. *J. Neurosci.* 23, 1087–1097.
- Beloozerova, I.N., Farrell, B.J., Sirota, M.G., and Prilutsky, B.I. (2010). Differences in movement mechanics, electromyographic, and motor cortex activity between accurate and nonaccurate stepping. *J. Neurophysiol.* 103, 2285–2300.
- Bennett, C., Arroyo, S., and Hestrin, S. (2013). Subthreshold mechanisms underlying state-dependent modulation of visual responses. *Neuron* 80, 350–357.
- Berridge, C.W., and Waterhouse, B.D. (2003). The locus coeruleus-noradrenergic system: modulation of behavioral state and state-dependent cognitive processes. *Brain Res. Brain Res. Rev.* 42, 33–84.
- Brozović, M., Abbott, L.F., and Andersen, R.A. (2008). Mechanism of gain modulation at single neuron and network levels. *J. Comput. Neurosci.* 25, 158–168.

- Carter, M.E., Yizhar, O., Chikahisa, S., Nguyen, H., Adamantidis, A., Nishino, S., Deisseroth, K., and de Lecea, L. (2010). Tuning arousal with optogenetic modulation of locus coeruleus neurons. *Nat. Neurosci.* **13**, 1526–1533.
- Castro-Alamancos, M.A. (2004). Absence of rapid sensory adaptation in neocortex during information processing states. *Neuron* **41**, 455–464.
- Castro-Alamancos, M.A., and Connors, B.W. (1996). Short-term plasticity of a thalamocortical pathway dynamically modulated by behavioral state. *Science* **272**, 274–277.
- Castro-Alamancos, M.A., and Gulati, T. (2014). Neuromodulators produce distinct activated states in neocortex. *J. Neurosci.* **34**, 12353–12367.
- Castro-Alamancos, M.A., and Oldford, E. (2002). Cortical sensory suppression during arousal is due to the activity-dependent depression of thalamocortical synapses. *J. Physiol.* **541**, 319–331.
- Chance, F.S., Abbott, L.F., and Reyes, A.D. (2002). Gain modulation from background synaptic input. *Neuron* **35**, 773–782.
- Constantinople, C.M., and Bruno, R.M. (2011). Effects and mechanisms of wakefulness on local cortical networks. *Neuron* **69**, 1061–1068.
- Constantinople, C.M., and Bruno, R.M. (2013). Deep cortical layers are activated directly by thalamus. *Science* **340**, 1591–1594.
- Costa, R.M., Cohen, D., and Nicolelis, M.A. (2004). Differential corticostriatal plasticity during fast and slow motor skill learning in mice. *Curr. Biol.* **14**, 1124–1134.
- Cowan, R.L., and Wilson, C.J. (1994). Spontaneous firing patterns and axonal projections of single corticostriatal neurons in the rat medial agranular cortex. *J. Neurophysiol.* **71**, 17–32.
- Crochet, S., and Petersen, C.C. (2006). Correlating whisker behavior with membrane potential in barrel cortex of awake mice. *Nat. Neurosci.* **9**, 608–610.
- Deuchars, J., West, D.C., and Thomson, A.M. (1994). Relationships between morphology and physiology of pyramid-pyramid single axon connections in rat neocortex *in vitro*. *J. Physiol.* **478**, 423–435.
- Dombeck, D.A., Graziano, M.S., and Tank, D.W. (2009). Functional clustering of neurons in motor cortex determined by cellular resolution imaging in awake behaving mice. *J. Neurosci.* **29**, 13751–13760.
- Eggermann, E., Kremer, Y., Crochet, S., and Petersen, C.C. (2014). Cholinergic signals in mouse barrel cortex during active whisker sensing. *Cell Rep.* **9**, 1654–1660.
- Favero, M., Varghese, G., and Castro-Alamancos, M.A. (2012). The state of somatosensory cortex during neuromodulation. *J. Neurophysiol.* **108**, 1010–1024.
- Forssberg, H., Grillner, S., Halbertsma, J., and Rossignol, S. (1980). The locomotion of the low spinal cat. II. Interlimb coordination. *Acta Physiol. Scand.* **108**, 283–295.
- Fu, Y., Tucciarone, J.M., Espinosa, J.S., Sheng, N., Darcy, D.P., Nicoll, R.A., Huang, Z.J., and Stryker, M.P. (2014). A cortical circuit for gain control by behavioral state. *Cell* **156**, 1139–1152.
- Grillner, S. (1981). Control of locomotion in bipeds, tetrapods and fish. In *Handbook of Physiology* (Bethesda: American Physiology Society), pp. 1179–1236.
- Grillner, S., and Zangger, P. (1979). On the central generation of locomotion in the low spinal cat. *Exp. Brain Res.* **34**, 241–261.
- Hattox, A.M., and Nelson, S.B. (2007). Layer V neurons in mouse cortex projecting to different targets have distinct physiological properties. *J. Neurophysiol.* **98**, 3330–3340.
- Hirata, A., and Castro-Alamancos, M.A. (2010). Neocortex network activation and deactivation states controlled by the thalamus. *J. Neurophysiol.* **103**, 1147–1157.
- Hô, N., and Destexhe, A. (2000). Synaptic background activity enhances the responsiveness of neocortical pyramidal neurons. *J. Neurophysiol.* **84**, 1488–1496.
- Hooks, B.M., Mao, T., Gutnisky, D.A., Yamawaki, N., Svoboda, K., and Shepherd, G.M. (2013). Organization of cortical and thalamic input to pyramidal neurons in mouse motor cortex. *J. Neurosci.* **33**, 748–760.
- Kaneko, T., Caria, M.A., and Asanuma, H. (1994). Information processing within the motor cortex. II. Intracortical connections between neurons receiving somatosensory cortical input and motor output neurons of the cortex. *J. Comp. Neurol.* **345**, 172–184.
- Kita, T., and Kita, H. (2012). The subthalamic nucleus is one of multiple innervation sites for long-range corticofugal axons: a single-axon tracing study in the rat. *J. Neurosci.* **32**, 5990–5999.
- Leone, D.P., Srinivasan, K., Chen, B., Alcamo, E., and McConnell, S.K. (2008). The determination of projection neuron identity in the developing cerebral cortex. *Curr. Opin. Neurobiol.* **18**, 28–35.
- Lorang, D., Amara, S.G., and Simerly, R.B. (1994). Cell-type-specific expression of catecholamine transporters in the rat brain. *J. Neurosci.* **14**, 4903–4914.
- Marlinski, V., Nilaweera, W.U., Zelenin, P.V., Sirota, M.G., and Beloozerova, I.N. (2012). Signals from the ventrolateral thalamus to the motor cortex during locomotion. *J. Neurophysiol.* **107**, 455–472.
- Murphy, B.K., and Miller, K.D. (2003). Multiplicative gain changes are induced by excitation or inhibition alone. *J. Neurosci.* **23**, 10040–10051.
- Orlovsky, G.N. (1972). The effect of different descending systems on flexor and extensor activity during locomotion. *Brain Res.* **40**, 359–371.
- Polack, P.O., Friedman, J., and Golshani, P. (2013). Cellular mechanisms of brain state-dependent gain modulation in visual cortex. *Nat. Neurosci.* **16**, 1331–1339.
- Poulet, J.F., Fernandez, L.M., Crochet, S., and Petersen, C.C. (2012). Thalamic control of cortical states. *Nat. Neurosci.* **15**, 370–372.
- Reyes, A., and Sakmann, B. (1999). Developmental switch in the short-term modification of unitary EPSPs evoked in layer 2/3 and layer 5 pyramidal neurons of rat neocortex. *J. Neurosci.* **19**, 3827–3835.
- Sheets, P.L., Suter, B.A., Kiritani, T., Chan, C.S., Surmeier, D.J., and Shepherd, G.M.G. (2011). Corticospinal-specific HCN expression in mouse motor cortex: I(h)-dependent synaptic integration as a candidate microcircuit mechanism involved in motor control. *J. Neurophysiol.* **106**, 2216–2231.
- Steriade, M., Amzica, F., and Nuñez, A. (1993a). Cholinergic and noradrenergic modulation of the slow (approximately 0.3 Hz) oscillation in neocortical cells. *J. Neurophysiol.* **70**, 1385–1400.
- Steriade, M., McCormick, D.A., and Sejnowski, T.J. (1993b). Thalamocortical oscillations in the sleeping and aroused brain. *Science* **262**, 679–685.
- Steriade, M., Nuñez, A., and Amzica, F. (1993c). A novel slow (< 1 Hz) oscillation of neocortical neurons in vivo: depolarizing and hyperpolarizing components. *J. Neurosci.* **13**, 3252–3265.
- Timofeev, I., Grenier, F., and Steriade, M. (2001). Disfacilitation and active inhibition in the neocortex during the natural sleep-wake cycle: an intracellular study. *Proc. Natl. Acad. Sci. USA* **98**, 1924–1929.
- Ueno, M., and Yamashita, T. (2011). Kinematic analyses reveal impaired locomotion following injury of the motor cortex in mice. *Exp. Neurol.* **230**, 280–290.
- Wang, Z., and McCormick, D.A. (1993). Control of firing mode of corticotectal and corticopontine layer V burst-generating neurons by norepinephrine, acetylcholine, and 1S,3R-ACPD. *J. Neurosci.* **13**, 2199–2216.
- Wang, M., Ramos, B.P., Paspalas, C.D., Shu, Y., Simen, A., Duque, A., Vijayraghavan, S., Brennan, A., Dudley, A., Nou, E., et al. (2007). Alpha2A-adrenoceptors strengthen working memory networks by inhibiting cAMP-HCN channel signaling in prefrontal cortex. *Cell* **129**, 397–410.
- Weiler, N., Wood, L., Yu, J., Solla, S.A., and Shepherd, G.M. (2008). Top-down laminar organization of the excitatory network in motor cortex. *Nat. Neurosci.* **11**, 360–366.